

Risk Factors Filtering

The 'Risks Factors' Filter allows practices to filter for patients with risk factors: Smoking, Alcohol (Drinker and High Risk), Drug Abuse, Pregnancy, Obesity and transplant patient. In addition it allows filtering for patients with conditions that are associated with an increased risk of pneumococcal for immunisation purposes. The risk factors are defined as:

- Smoking – daily or irregular smokers
- Alcohol – drinker: drinks alcohol
- Alcohol – drinker high risk: 2 or more drinks on a regular occasion or more than 4 drinks on any occasion
- Medication/Drug abuse – coded diagnosis in patient record
- Pneumococcal – has a condition that requires immunisation is up to date – please refer to the page about adult immunisation for full details. This page can be found here: [Details of Risk Factor Filter Pneumococcal](#)
- Pregnancy – has active pregnancy in clinical record
- Obesity – BMI of 30 or more
- Transplant - patient has received an organ transplant. For mapping details please refer to the conditions mapping for your clinical system on our [Data Mapping](#) pages
- Fracture - patients with a record of one or more fractures. For mapping details please refer to the conditions mapping for your clinical system on our [Data Mapping](#) pages

Filter

General

Ethnicity

Conditions

Medications

Date Range (Results)

Date Range (Visits)

Patient Name

Patient Status

Providers

Risk Factors

Health Care Homes

MBS Attendance

Custom Filters

Saved Filters

Risk Factors

Substance Abuse

☐ Smoking

☐ Alcohol - Drinker

☐ Alcohol - Drinker High Risk *

☐ Medication/Drug abuse **

Requires Immunisation

Has Conditions increasing risk of

☐ Pneumococcal

Other Risk Categories

☐ Pregnancy

☐ Obesity

☐ Transplant

☐ Fracture

* 2 or more drinks on a regular occasion or more than 4 drinks on any occasion ** Diagnosis

Clear